

On the Sudakov suppression and enhancement for electroweak reactions

B.I. Ermolaev¹

¹*CFTC, University of Lisbon Av. Prof. Gama Pinto 2, P-1649-003 Lisbon, Portugal
and*

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

Accounting for the double-logarithmic (DL) contributions to amplitude of $Z \rightarrow f \bar{f}$ leads to the Sudakov suppression. In contrast, resumming DL contributions to amplitudes of $W^\pm \rightarrow f \bar{f}'$ results into an exponential with a positive exponent, i.e. into the enhancement. This effect is intrinsic property of the theories with non-Abelian gauge groups.

PACS numbers: 12.38.Cy

INTRODUCTION

Among different double-logarithmic (DL) contributions, there are the contributions that depend on the infrared cut-off. Results of resumming them to all orders in the coupling(s) can be often written in an exponential form, with a negative exponent, so that it falls when the total energy grows. This is known as the Sudakov suppression [1]. It means the suppression of cross-sections of non-radiative processes compared to cross sections of the processes where bremsstrahlung is allowed. Such a suppression in QED (see e.g review [2]) and in QCD[3] was studied in details to all orders in the couplings. Exponentiation of the first-loop electroweak (EW) contributions was obtained in Ref. [4] and then was confirmed in Refs. [5]-[9] by the fixed orders calculations. The complete list of publications on this subject can be found in the recent Refs. [10],[11].

However, resumming DL contributions of the same infrared origin sometimes leads to different results. In particular, it was found in Ref. [12] that one-loop DL contribution to the $W^\pm \rightarrow f \bar{f}'$ has the opposite sign compared to the one for $\gamma \rightarrow f \bar{f}$ and $Z \rightarrow f \bar{f}$ -decays. It suggests that the total resummation of the DL corrections to the amplitude of the $W^\pm \rightarrow f \bar{f}'$ may be an exponential with a positive exponent.

In the present paper we consider amplitudes for the W -decay. We show that the result of accounting for DL contributions to all orders in the EW couplings has an exponential form with the positive exponent. In other words, there is the enhancement instead of the suppression for such amplitudes. We consider the decays of the electroweak bosons into the left fermions f and the right antifermions $\bar{f}$, $\bar{f}'$ though drop the subscripts L and R through the paper.

The paper is organised as follows: in Sect. II we briefly review the Sudakov suppression for amplitude of $Z \rightarrow f \bar{f}$, comparing the use the Feynman and the Coulomb gauges. In Sect. III we discuss the decay of W -bosons into fermions, again with using both the Feynman and

the Coulomb gauges, and arrive at the enhancement. The origin of the enhancement is considered in Sect. IV. Finally, Sect. V is for concluding remarks.

SUDAKOV SUPPRESSION

The Sudakov suppression[1] is well-known, so we discuss it very briefly. Let us consider amplitude A_Z of the decay $Z \rightarrow f \bar{f}$ where f stands for any (charged) fermion from the left doublets of the Standard Model. As we assume all external particles to be on-shell, A_Z is gauge-independent. Among the often used gauges for propagators $D_{\mu\nu}(k)$ of the EW bosons, there are the Feynman gauge and the Coulomb (temporal) gauge.




FIG. 1:

In the first-loop approximation and when the Feynman gauge is used, the DL contribution to A_Z comes from the graph in Fig. 1. The vertical wavy line in Fig. 1 denotes the external EW boson. The horizontal wavy line may correspond to any virtual electroweak boson though the DL contribution to A_Z appears only when this boson is a photon. When the Coulomb (temporal) gauge is used, DL contributions can come from the self-energy graphs depicted in Fig. 2. Again, DL contributions come only when the virtual bosons in Fig. 2 are photons.

Using the Feynman gauge, it is easy to arrive at the well-known one-loop DL contribution $A_Z^{(1)}$ to A_Z :

$$A_Z^{(1)}/A_Z^{Born} = -Q_f^2 \left(\frac{\alpha}{4\pi} \ln^2(-M^2/\mu^2) \right) \equiv -Q_f^2 L \quad (1)$$




FIG. 2:

where Q_f is the electric charge of the fermion f , M is the mass of the Z -boson and μ is the infrared cut-off. With DL accuracy, one can neglect the difference between masses of Z and W -bosons. The double logarithm in Eq. (1) is written in the simplest way for the particular case when $\mu \geq m$, with m being the mass of the fermion f . It is also easy to arrive at Eq. (1) when the Coulomb gauge is used. In this case, DL contributions come from the self-energy graphs depicted in Fig. 2. The DL contributions Σ_f^{DL} from the fermion self-energy graphs (graphs (b) and (c) in Fig. 2) are $\Sigma_f^{DL} = -Q_f^2 L/2$. In order to get a DL contribution from graph (a) in Fig. 2, at least one of the virtual bosons in Fig. 2a has to be a photon. However, there is not such a vertex in the Lagrangian of the Standard Model. It leaves us with two contributions Σ_f^{DL} and we are back to Eq. (1). Using the Coulomb gauge for the Sudakov logarithms was discussed in details in Refs. [13],[9]. The DL asymptotics of A_Z accounting for contributions both $\sim \gamma_\mu$ and $\sim \sigma_{\mu\nu} = (1/2)(\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu)$ can be obtained by direct graph-by-graph calculations like it was done in Ref. [14]. The complete expression for the DL asymptotics for A_Z is

$$A_Z = \left[\gamma_\mu + \frac{\sigma_{\mu\nu} q_\nu}{m} \frac{\partial}{\partial \rho} \right] e^{-Q_f^2 (\alpha/4\pi) \ln^2 \rho} \quad (2)$$

where we have omitted the spinors of the fermions and the Z -boson polarization vector, q is the sum of the fermion momenta ($q^2 = M^2$); we have chosen in Eq. (2) $\mu \approx m$ and denoted $\rho \equiv -M^2/m^2$. The minus sign in the exponential of Eq. (2) means the Sudakov suppression. It suppresses the non-radiative decays of Z -bosons.

SUDAKOV ENHANCEMENT

Now let us consider DL contributions to amplitude A_W of the decay $W^\pm \rightarrow f \bar{f}'$ where f and f' ($f \neq f'$) belong to the same doublet of the Standard Model. When the Feynman gauge is used, DL contribution comes from the graph in Fig. 1 where the horizontal wavy line denotes again the virtual photon. This contribution is

$$A_W^{(1)}/A_W^{(Born)} = -Q_f Q_{f'} L \quad (3)$$

where $Q_f(Q_{f'})$ is the electric charge of the fermion f (f'). Invoking results of Ref. [14] leads immediately to the DL asymptotics for A_W :

$$A_W = \left[\gamma_\mu + \frac{\sigma_{\mu\nu} q_\nu}{m} \frac{\partial}{\partial \rho} \right] e^{-Q_f Q_{f'} (\alpha/4\pi) \ln^2 \rho}. \quad (4)$$

The term $\sim \alpha$ in the expansion of Eq. (4) into series in α was obtained in Ref. [12].

Let us notice that when the fermions f and f' in Eq. (4) are quarks, they have opposite signs. It means that the exponent in Eq. (4) is positive and therefore there is the DL enhancement in Eq. (4) instead of the DL suppression of Eq. (2). For reactions $W^+ \rightarrow u \bar{d}$ and $W^- \rightarrow d \bar{u}$, the DL amplitude of the process is

$$A_W(\text{quarks}) = A_W^{(0)}(\text{quarks}) e^{(2/9)L}. \quad (5)$$

Notation $A_W^{(0)}(\text{quarks})$ in Eq. (5) stands for the expression in the squared brackets in Eq. (4). At the same time, Eq. (4) reads that there are no DL contributions to the lepton decays $W \rightarrow l \bar{\nu}_l$, $W \rightarrow \nu_l \bar{l}$. Although amplitudes A_W are gauge-invariant, it is useful to demonstrate how the first-loop contribution $A_W^{(1)}$ of Eq. (3) can be easily obtained when the Coulomb gauge is used. In this case, DL contributions come from the self-energy graphs. Contributions of graphs (b) and (c) in Fig. 2 are $\Sigma_f^{DL} = -Q_f^2 L/2$ and $\Sigma_{f'}^{DL} = -Q_{f'}^2 L/2$ respectively. Contrary to the case of the Z -boson decay, graph (a) in Fig. 1 now yields a DL contribution

$$P_W^{DL} = Q_W^2 L/2 \quad (6)$$

and therefore

$$A_W^{(1)} = A_W^{Born} \left[P_W^{DL} + \Sigma_f^{DL} + \Sigma_{f'}^{DL} \right] = A_W^{Born} [Q_W^2 - Q_f^2 - Q_{f'}^2] L/2. \quad (7)$$

The electric charge conservation states that $Q_f - Q_{f'} = Q_W$. It allows to rewrite Eq. (7) in the form of Eq. (3).

ORIGIN OF THE ENHANCEMENT

Both the Sudakov suppression of Eq. (2) and the enhancement of Eq. (4) have a very simple origin in terms of the algebra of the gauge group generators. We explain it, first using the Feynman gauge and then proceed to the case of the Coulomb gauge. To begin with, let us remind that all electroweak boson fields A , Z , W are linear combinations of the gauge isoscalar field B and isovector fields A_i ($i = 1, 2, 3$):

$$W^\pm = \frac{A_1 \pm iA_2}{\sqrt{2}}, \quad Z = -B \sin \theta_W + A_3 \cos \theta_W, \\ A = B \cos \theta_W + A_3 \sin \theta_W \quad (8)$$

where θ_W is the Weinberg angle. Therefore, propagator $D_{\mu\nu}^{(B)}$ of the isoscalar field B and propagator $D_{\mu\nu}^{(3)}$ of the isovector field A_3 can be expressed in terms of the photon propagator $D_{\mu\nu}^{(A)}$ and the Z -boson propagator $D_{\mu\nu}^{(Z)}$. As DL contributions at the energies $\leq M$ come from the photon propagators only,

$$D_{\mu\nu}^{(B)} \approx D_{\mu\nu}^{(A)} \cos^2 \theta_W, \quad D_{\mu\nu}^{(3)} \approx D_{\mu\nu}^{(A)} \sin^2 \theta_W. \quad (9)$$

The $SU(2)U(1)$ -parts of the vertexes of the interaction of fields A_i and B with fermions are $gt_i A_i$ and $g'(Y/2)B$, with t_i being the $SU(2)$ -generators ($i = 1, 2, 3$), Y being the fermion hypercharge and g, g' being the electroweak couplings ($g' = g \tan \theta_W$). Now let us choose the Feynman gauge for the photon propagator and consider again the one-loop graph in Fig. 1. We can regard both the external and virtual electroweak bosons in this Fig. as consisting of the gauge fields B, A_i . In order to discuss both the Z and the W -decays at the same time, let us suppose that the external field X_r consists of the isoscalar and isovector fields: $X_r = pB + qA_r$ where $r = 1, 2, 3$; p and q are some scalar factors. Then the Born amplitude can be written as $p + qt_r$ and the graph in Fig. 1 yields the following DL contribution:

$$A_r^{(1)} = - \left[g^2 \sin^2 \theta_W (t_3(p + qt_r)t_3) + \right. \\ \left. g'^2 \cos^2 \theta_W \frac{Y^2}{4} (pB + qt_r) \right] (1/16\pi^2) \ln^2(-M^2/\mu^2). \quad (10)$$

Therefore,

$$A_r^{(1)} = -A_r^{(Born)} \frac{(\xi_r + Y^2)}{4} L \quad (11)$$

where $\xi_3 = 1$, $\xi_{1,2} = -1$

Eq. (11) leads to the suppression when $r = 3$ and to the enhancement when $r = 1, 2$ provided $Y^2 < 1$. The exponentiation of DL contributions in terms of the

gauge fields B, A_3 can be proved easily, e.g. by composing the Infrared Evolution Equations for amplitudes A_r in the same way as it was done in Ref. [14] for calculating the DL asymptotics of the form factors of electron (Abelian gauge group) and quark (non-Abelian gauge group). Eq. (11) reads that the first-loop DL contributions from exchange of the virtual isoscalar boson (the contributions $\sim Y^2$) always lead to the suppression whereas the exchanges of the isovector bosons lead either to the suppression or to the enhancement due to non-commutativity of the generators of the gauge group.

In order to obtain this result when the Coulomb gauge is used, it is enough to notice that the only difference between DL contributions of the graphs in Fig. 2 compared to Eqs. (6,7) is in $SU(2)$ -group factors. After accounting for them, one reproduces Eq. (11).

SUMMARY AND OUTLOOK

We have considered the decays of the massive electroweak bosons into fermions in the DL approximation. We have shown that when the EW bosons are charged, accounting for the DL radiative corrections leads to exponentiation of the first-loop contribution, however with the positive exponent, i.e. instead of the Sudakov suppression there is the enhancement for such decays. On the other hand, accounting for the soft photon emission accompanying such decays leads to the falling exponentials. Therefore, these DL contributions cancel each other in expressions for inclusive cross-sections. The results we obtained can be useful for analyses of $1 \rightarrow 2$ processes in theories with spontaneously broken non-Abelian gauge groups which involve different mass scales for charged massive bosons.

ACKNOWLEDGEMENT

The work is supported by grant CERN/FIS/43652/2001.

-
- [1] V.V. Sudakov. ZhETP 30(1956)87.
 - [2] V.G. Gorshkov. Uspekhi Fiz. Nauk, 110(1973)45.
 - [3] J.J. Carazzone, E.C. Poggio and H.R. Quinn. Phys. Rev. D11(1975)2286; J.M. Cornwall and G. Tiktopoulos. Phys. Rev. Lett. 35(1975)338; Phys. Rev. D13(1976)3370; J. Frenkel and J.C.Taylor. Nucl. Phys. B116(1976)185; V.V. Belokurov and N.I. Usyukina. Phys. Rev. Lett. B94(1980)251; Theor. Math. Phys. 44(1980)657; ibid 45(1980)957.
 - [4] V.S. Fadin, L.N. Lipatov, A.D. Martin, and M. Melles. Phys.Rev. D61 (2000) 094002.

- [5] J.H. Kuhn, A.A. Penin and V.A. Smirnov. Nucl.Phys.Proc.Suppl. 89(2000)94; Eur. Phys. J.C17(2000)97.
- [6] M. Beccaria, P. Ciafaloni, D. Comeli, F. Renard, C. Verzegnassi. Phys. Rev. D 61(2000)073005 .
- [7] M. Melles Phys. Lett. B 495(2000)81 .
- [8] M. Hori, H. Kawamura and J. Kodaira. Phys. Lett. B 491(2000)275.
- [9] W. Beenakker and A. Werthenbach. Nucl. Phys. B 630(2002)3.
- [10] U. Baur and D. Wackeroth. hep-ph/0211089.
- [11] S. Moch. hep-ph/0211193.
- [12] D. Wackeroth and W. Hollik. Phys. Rev. D 55(1997)6788.
- [13] A.V. Smilga and M.I. Vysotsky. JETP Lett. 29(1979)251.
- [14] B.I. Ermolaev and S.I. Troyan. Nucl. Phys. B 590(2000)521 (hep-ph/0009345).